

## 4. Measurement of Angles

### Exercise 4.1

#### 1. Question

Find the degree measure corresponding to the following radian measures (Use  $\pi = \frac{22}{7}$ )

(i)  $\frac{9\pi}{5}$  (ii)  $-\frac{5\pi}{6}$  (iii)  $\left(\frac{18\pi}{5}\right)^c$  (iv)  $(-3)^c$  (v)  $11^c$  (vi)  $1^c$

#### Answer

We know that  $\pi \text{ rad} = 180^\circ \Rightarrow 1 \text{ rad} = 180^\circ / \pi$

(i) Given  $\frac{9\pi}{5}$

$$= \left(\frac{180}{\pi} \times \frac{9\pi}{5}\right)^\circ$$

$$= (36 \times 9)^\circ$$

$$= 324^\circ$$

(ii) Given  $-\frac{5\pi}{6}$

$$= \left(\frac{180}{\pi} \times -\frac{5\pi}{6}\right)^\circ$$

$$= (30 \times -5)^\circ$$

$$= -(150)^\circ$$

(iii) Given  $\left(\frac{18\pi}{5}\right)^c$

$$= \left(\frac{180}{\pi} \times \frac{18\pi}{5}\right)^\circ$$

$$= (36 \times 18)^\circ$$

$$= 648^\circ$$

(iv) Given  $(-3)^c$

$$= \left(\frac{180}{\pi} \times -3\right)^\circ$$

$$= \left(\frac{180}{22} \times 7 \times -3\right)^\circ$$

$$= \left(-\frac{3780}{22}\right)^\circ$$

$$= \left(-171\frac{18}{22}\right)^\circ$$

$$= \left(-171^\circ \left(\frac{18}{22} \times 60\right)'\right)$$

$$= \left(-171^\circ \left(49\frac{1}{11}\right)'\right)$$

Bodhiyla.com

$$= \left( -171^\circ 49' \left( \frac{1}{11} \times 60 \right)' \right)$$

$$= -(171^\circ 49' 5.45'')$$

$$\approx -(171^\circ 49' 5'')$$

(v) Given  $11^\circ$

$$= \left( \frac{180}{\pi} \times 11 \right)^\circ$$

$$= \left( \frac{180}{22} \times 7 \times 11 \right)^\circ$$

$$= (90 \times 7)^\circ$$

$$= 630^\circ$$

(vi) Given  $1^\circ$

$$= \left( \frac{180}{\pi} \times 1 \right)^\circ$$

$$= \left( \frac{180}{22} \times 7 \times 1 \right)^\circ$$

$$= \left( \frac{1260}{22} \right)^\circ$$

$$= \left( 57 \frac{3}{11} \right)^\circ$$

$$= \left( 57^\circ \left( \frac{3}{11} \times 60 \right)' \right)$$

$$= \left( 57^\circ \left( 16 \frac{4}{11} \right)' \right)$$

$$= \left( 57^\circ 16' \left( \frac{4}{11} \times 60 \right)' \right)$$

$$= (57^\circ 16' 21.81'')$$

$$\approx (57^\circ 16' 22'')$$

Bodhiyla.com

## 2. Question

Find the radian measure corresponding to the following degree measures:

(i)  $300^\circ$  (ii)  $35^\circ$  (iii)  $-56^\circ$  (iv)  $135^\circ$  (v)  $-300^\circ$

(vi)  $7^\circ 30'$  (vii)  $125^\circ 30'$  (viii)  $-47^\circ 30'$

## Answer

We know that  $180^\circ = \pi \text{ rad} \Rightarrow 1^\circ = \pi/180 \text{ rad}$

(i) Given  $300^\circ$

$$= \left( 300 \times \frac{\pi}{180} \right) \text{ rad}$$

$$= \frac{5\pi}{3} \text{ rad}$$

(ii) Given  $35^\circ$

$$= \left( 35 \times \frac{\pi}{180} \right) \text{ rad}$$

$$= \frac{7\pi}{36} \text{ rad}$$

(iii) Given  $-56^\circ$

$$= \left( -56 \times \frac{\pi}{180} \right) \text{ rad}$$

$$= -\frac{14\pi}{45} \text{ rad}$$

(iv) Given  $135^\circ$

$$= \left( 135 \times \frac{\pi}{180} \right) \text{ rad}$$

$$= \frac{3\pi}{4} \text{ rad}$$

(v) Given  $-300^\circ$

$$= \left( -300 \times \frac{\pi}{180} \right) \text{ rad}$$

$$= -\frac{5\pi}{3} \text{ rad}$$

(vi) Given  $7^\circ 30'$

We know that  $30' = (1/2)^\circ$

$$\therefore 7^\circ 30' = (7 \frac{1}{2})^\circ$$

$$= \left( \frac{15}{2} \right)^\circ$$

$$= \left( \frac{15}{2} \times \frac{\pi}{180} \right) \text{ rad}$$

$$= \left( \frac{\pi}{24} \right) \text{ rad}$$

(vii) Given  $125^\circ 30'$

We know that  $30' = (1/2)^\circ$

$$\therefore 125^\circ 30' = (125 \frac{1}{2})^\circ$$

$$= \left( \frac{251}{2} \right)^\circ$$

$$= \left( \frac{251}{2} \times \frac{\pi}{180} \right) \text{ rad}$$

$$= \left( \frac{251\pi}{360} \right) \text{ rad}$$

(viii) Given  $-47^\circ 30'$

We know that  $30' = (1/2)^\circ$

$$\therefore -47^\circ 30' = -(47 \frac{1}{2})^\circ$$

$$= -\left( \frac{95}{2} \right)^\circ$$

Bodhiyla.com

$$= -\left(\frac{95}{2} \times \frac{\pi}{180}\right) \text{rad}$$

$$= -\left(\frac{19\pi}{72}\right) \text{rad}$$

### 3. Question

The difference between the two acute angles of a right-angled triangle is  $2\pi/5$  radians. Express the angles in degrees.

### Answer

Given the difference between the two acute angles of a right-angled triangle is  $2\pi/5$  radians.

We know that  $\pi \text{ rad} = 180^\circ \Rightarrow 1 \text{ rad} = 180^\circ/\pi$

Given  $\frac{2\pi}{5}$

$$= \left(\frac{180}{\pi} \times \frac{2\pi}{5}\right)^\circ$$

$$= (36 \times 2)^\circ$$

$$= 72^\circ$$

Let one acute angle be  $x^\circ$  and the other acute angle be  $90^\circ - x^\circ$ .

Then,

$$\Rightarrow x^\circ - (90^\circ - x^\circ) = 72^\circ$$

$$\Rightarrow 2x^\circ - 90^\circ = 72^\circ$$

$$\Rightarrow 2x^\circ = 72^\circ + 90^\circ$$

$$\Rightarrow 2x^\circ = 162^\circ$$

$$\Rightarrow x^\circ = 162^\circ/2$$

$$\therefore x^\circ = 81^\circ \text{ and } 90^\circ - x^\circ = 90^\circ - 81^\circ = 9^\circ$$

### 4. Question

One angle of a triangle is  $\frac{2}{3}x$  grades, and another is  $\frac{3}{2}x$  degrees while the third is  $\frac{\pi x}{75}$  radians. Express all the angles in degrees.

### Answer

Given one angle of a triangle is  $2x/3$  grades and another is  $3x/2$  degree while the third is  $\pi x/75$  radians.

We know that  $1 \text{ grad} = \left(\frac{9}{10}\right)^\circ$

$$\Rightarrow \frac{2}{3}x \text{ grad} = \left(\frac{9}{10}\right)\left(\frac{2}{3}x\right)^\circ = \frac{3}{5}x^\circ$$

We know that  $\pi \text{ rad} = 180^\circ \Rightarrow 1 \text{ rad} = 180^\circ/\pi$

Given  $\frac{\pi x}{75}$

$$= \left(\frac{180}{\pi} \times \frac{\pi x}{75}\right)^\circ$$

$$= \left(\frac{12}{5}x\right)^\circ$$

We know that the sum of the angles of a triangle is  $180^\circ$ .

$$\Rightarrow \frac{3}{5}x^\circ + \frac{3}{2}x^\circ + \frac{12}{5}x^\circ = 180^\circ$$

$$\Rightarrow \frac{6 + 15 + 24}{10}x^\circ = 180^\circ$$

$$\Rightarrow 45x^\circ = 180^\circ \times 10^\circ$$

$$\Rightarrow 45x^\circ = 1800^\circ$$

$$\Rightarrow x^\circ = 1800^\circ / 45^\circ$$

$$\therefore x = 40^\circ$$

$\therefore$  The angles of the triangle are

$$\Rightarrow \frac{3}{5}x^\circ = \frac{3}{5} \times 40^\circ = 24^\circ$$

$$\Rightarrow \frac{3}{2}x^\circ = \frac{3}{2} \times 40^\circ = 60^\circ$$

$$\Rightarrow \frac{12}{5}x^\circ = \frac{12}{5} \times 40^\circ = 96^\circ$$

### 5. Question

Find the magnitude, in radians and degrees, of the interior angle of a regular:

(i) Pentagon (ii) Octagon (iii) Heptagon (iv) Duodecagon.

### Answer

We know that the sum of the interior angles of a polygon =  $(n - 2)\pi$

And each angle of polygon =  $\frac{\text{sum of interior angles of polygon}}{\text{number of sides}}$

(i) Pentagon

Number of sides in pentagon = 5

Sum of interior angles of pentagon =  $(5 - 2)\pi = 3\pi$

$$\therefore \text{Each angle of pentagon} = \frac{3\pi}{5} \times \frac{180^\circ}{\pi} = 108^\circ$$

(ii) Octagon

Number of sides in octagon = 8

Sum of interior angles of octagon =  $(8 - 2)\pi = 6\pi$

$$\therefore \text{Each angle of octagon} = \frac{6\pi}{8} \times \frac{180^\circ}{\pi} = 135^\circ$$

(iii) Heptagon

Number of sides in heptagon = 7

Sum of interior angles of heptagon =  $(7 - 2)\pi = 5\pi$

$$\therefore \text{Each angle of heptagon} = \frac{5\pi}{7} \times \frac{180^\circ}{\pi} = \frac{900}{7}^\circ = 128^\circ 34' 17''$$

(iv) Duodecagon

Number of sides in duodecagon = 12

Sum of interior angles of duodecagon =  $(12 - 2)\pi = 10\pi$

$$\therefore \text{Each angle of duodecagon} = \frac{10\pi}{12} \times \frac{180^\circ}{\pi} = 150^\circ$$

## 6. Question

The angles of a quadrilateral are in A.P., and the greatest angle is  $120^{\circ}$ . Express the angles in radians.

### Answer

Let the angles of quadrilateral be  $(a - 3d)^{\circ}$ ,  $(a - d)^{\circ}$ ,  $(a + d)^{\circ}$  and  $(a + 3d)^{\circ}$ .

We know that the sum of angles of a quadrilateral is  $360^{\circ}$ .

$$\Rightarrow a - 3d + a - d + a + d + a + 3d = 360^{\circ}$$

$$\Rightarrow 4a = 360^{\circ}$$

$$\therefore a = 90^{\circ}$$

Given the greatest angle =  $120^{\circ}$

$$\Rightarrow a + 3d = 120^{\circ}$$

$$\Rightarrow 90^{\circ} + 3d = 120^{\circ}$$

$$\Rightarrow 3d = 120^{\circ} - 90^{\circ}$$

$$\Rightarrow 3d = 30^{\circ}$$

$$\Rightarrow d = 10^{\circ}$$

Hence, the angles are:

$$\Rightarrow (a - 3d)^{\circ} = 90^{\circ} - 30^{\circ} = 60^{\circ}$$

$$\Rightarrow (a - d)^{\circ} = 90^{\circ} - 10^{\circ} = 80^{\circ}$$

$$\Rightarrow (a + d)^{\circ} = 90^{\circ} + 10^{\circ} = 100^{\circ}$$

$$\Rightarrow (a + 3d)^{\circ} = 120^{\circ}$$

Angles of quadrilateral in radians:

$$\Rightarrow \left(60 \times \frac{\pi}{180}\right) \text{ rad} = \frac{\pi}{3} \text{ rad}$$

$$\Rightarrow \left(80 \times \frac{\pi}{180}\right) \text{ rad} = \frac{4\pi}{9} \text{ rad}$$

$$\Rightarrow \left(100 \times \frac{\pi}{180}\right) \text{ rad} = \frac{5\pi}{9} \text{ rad}$$

$$\Rightarrow \left(120 \times \frac{\pi}{180}\right) \text{ rad} = \frac{2\pi}{3} \text{ rad}$$

## 7. Question

The angles of a triangle are in A.P., and the number of degrees in the least angle is to the number of degrees in the mean angle as 1:120. Find the angle in radians.

### Answer

Let the angles of the triangle be  $(a - d)^{\circ}$ ,  $a^{\circ}$  and  $(a + d)^{\circ}$ .

We know that the sum of the angles of a triangle is  $180^{\circ}$ .

$$\Rightarrow a - d + a + a + d = 180^{\circ}$$

$$\Rightarrow 3a = 180^{\circ}$$

$$\therefore a = 60^{\circ}$$

$$\text{Given } \frac{\text{Number of degrees in the least angle}}{\text{Number of degrees in the mean angle}} = \frac{1}{120}$$

$$\Rightarrow \frac{a-d}{a} = \frac{1}{120}$$

$$\Rightarrow \frac{60-d}{60} = \frac{1}{120}$$

$$\Rightarrow \frac{60-d}{1} = \frac{1}{2}$$

$$\Rightarrow 120 - 2d = 1$$

$$\Rightarrow 2d = 119$$

$$\therefore d = 59.5$$

Hence, angles are:

$$\Rightarrow (a-d)^\circ = 60^\circ - 59.5^\circ = 0.5^\circ$$

$$\Rightarrow a^\circ = 60^\circ$$

$$\Rightarrow (a+d)^\circ = 60^\circ + 59.5^\circ = 119.5^\circ$$

$\therefore$  Angles of triangle in radians:

$$\Rightarrow \left(0.5 \times \frac{\pi}{180}\right) \text{ rad} = \frac{\pi}{360} \text{ rad}$$

$$\Rightarrow \left(60 \times \frac{\pi}{180}\right) \text{ rad} = \frac{\pi}{3} \text{ rad}$$

$$\Rightarrow \left(119.5 \times \frac{\pi}{180}\right) \text{ rad} = \frac{239\pi}{360} \text{ rad}$$

### 8. Question

The angle in one regular polygon is to that in another as 3:2 and the number of sides in first is twice that in the second. Determine the number of sides of two polygons.

### Answer

Let the number of sides in the first polygon be  $2x$  and the number of sides in the second polygon be  $x$ .

We know that angle of an  $n$ -sided regular polygon =  $\left(\frac{n-2}{n}\right)\pi$  radian

$$\Rightarrow \text{The angle of the first polygon} = \left(\frac{2x-2}{2x}\right)\pi = \left(\frac{x-1}{x}\right)\pi \text{ radian}$$

$$\Rightarrow \text{The angle of the second polygon} = \left(\frac{x-2}{x}\right)\pi \text{ radian}$$

Thus,

$$\Rightarrow \frac{\left(\frac{x-1}{x}\right)\pi}{\left(\frac{x-2}{x}\right)\pi} = \frac{3}{2}$$

$$\Rightarrow \frac{x-1}{x-2} = \frac{3}{2}$$

$$\Rightarrow 2x - 2 = 3x - 6$$

$$\therefore x = 4$$

Thus,

Number of sides in the first polygon =  $2x = 8$

Number of sides in the second polygon =  $x = 4$

### 9. Question

The angles of a triangle are in A.P. such that the greatest is 5 times the least. Find the angles in radians.

### Answer

Let the angles of the triangle be  $(a - d)^\circ$ ,  $a^\circ$  and  $(a + d)^\circ$ .

We know that the sum of angles of triangle is  $180^\circ$ .

$$\Rightarrow a - d + a + a + d = 180^\circ$$

$$\Rightarrow 3a = 180^\circ$$

$$\therefore a = 60^\circ$$

Given greatest angle = 5 × least angle

$$\frac{\text{Greatest angle}}{\text{Least angle}} = 5$$

$$\Rightarrow \frac{a + d}{a - d} = 5$$

$$\Rightarrow \frac{60 + d}{60 - d} = 5$$

$$\Rightarrow 60 + d = 300 - 5d$$

$$\Rightarrow 6d = 240$$

$$\therefore d = 40$$

Hence, angles are:

$$\Rightarrow (a - d)^\circ = 60^\circ - 40^\circ = 20^\circ$$

$$\Rightarrow a^\circ = 60^\circ$$

$$\Rightarrow (a + d)^\circ = 60^\circ + 40^\circ = 100^\circ$$

$\therefore$  Angles of triangle in radians:

$$\Rightarrow \left(20 \times \frac{\pi}{180}\right) \text{ rad} = \frac{\pi}{9} \text{ rad}$$

$$\Rightarrow \left(60 \times \frac{\pi}{180}\right) \text{ rad} = \frac{\pi}{3} \text{ rad}$$

$$\Rightarrow \left(100 \times \frac{\pi}{180}\right) \text{ rad} = \frac{5\pi}{9} \text{ rad}$$

### 10. Question

The number of sides of two regular polygons is 5:4 and the difference between their angle is  $9^\circ$ . Find the number of sides of the polygons.

### Answer

Let the number of sides in the first polygon be  $5x$  and the number of sides in the second polygon be  $4x$ .

We know that angle of an  $n$ -sided regular polygon =  $\left(\frac{n-2}{n}\right) \pi$  radian

$$\Rightarrow \text{The angle of the first polygon} = \left(\frac{5x-2}{5x}\right) 180^\circ$$

$$\Rightarrow \text{The angle of the second polygon} = \left(\frac{4x-2}{4x}\right) 180^\circ$$

Thus,



$$\Rightarrow \left(\frac{5x-2}{5x}\right)180^\circ - \left(\frac{4x-1}{4x}\right)180^\circ = 9$$

$$\Rightarrow 180^\circ \left(\frac{4(5x-2) - 5(4x-1)}{20x}\right) = 9$$

$$\Rightarrow \frac{20x - 8 - 20x + 10}{20x} = \frac{9}{180}$$

$$\Rightarrow \frac{2}{20x} = \frac{1}{20}$$

$$\Rightarrow \frac{2}{x} = 1$$

$$\therefore x = 2$$

Thus,

Number of sides in the first polygon =  $5x = 10$

Number of sides in the second polygon =  $4x = 8$

### 11. Question

A railroad curve is to be laid out on a circle. What radius should be used if the track is to change direction by  $25^\circ$  at a distance of 40 meters?

### Answer

Given length of arc = 40 m

And  $\theta = 25^\circ$

We know that  $180^\circ = \pi \text{ rad} \Rightarrow 1^\circ = \pi/180 \text{ rad}$

$$\Rightarrow 25^\circ = \left(25 \times \frac{\pi}{180}\right) = \frac{5\pi}{36} \text{ radian}$$

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \frac{5\pi}{36} = \frac{40}{\text{radius}}$$

$$\Rightarrow \text{Radius} = \frac{40}{\frac{5\pi}{36}}$$

$$= \frac{40 \times 36 \times 7}{5 \times 22}$$

$$= 91.64 \text{ m}$$

So, the radius of the track should be 91.64 m.

### 12. Question

Find the length which at a distance of 5280 m will subtend an angle of  $1'$  at the eye.

### Answer

Given radius = 5280 m

We know that  $\theta = 1' = \left(\frac{1}{60}\right)^\circ = \left(\frac{1}{60} \times \frac{\pi}{180}\right) \text{ rad}$

And know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \frac{1}{60} \times \frac{\pi}{180} = \frac{\text{arc}}{5280}$$

$$\therefore \text{arc} = \frac{5280 \times 22}{60 \times 180 \times 7}$$

$$= 1.5365 \text{ m}$$

### 13. Question

A wheel makes 360 revolutions per minute. Through how many radians does it turn in 1 second?

### Answer

Given the number of revolutions taken by the wheel in 1 minute = 360

Number of revolution taken by the wheel in 1 second =  $360/6 = 6$

We know that 1 revolution =  $2\pi$  radians

$\therefore$  Number of radians the wheel will turn in 1 second =  $6 \times 2\pi = 12\pi$

### 14. Question

Find the angle in radians through which a pendulum swings if its length is 75 cm and the tip describes an arc of length (i) 10 cm (ii) 15 cm (iii) 21 cm.

### Answer

Given radius = 75 cm

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

(i) Given length of arc = 10 cm

$$\Rightarrow \theta = \frac{10}{75} = \frac{2}{15} \text{ radian}$$

(ii) Given length of arc = 15 cm

$$\Rightarrow \theta = \frac{15}{75} = \frac{1}{5} \text{ radian}$$

(iii) Given length of arc = 21 cm

$$\Rightarrow \theta = \frac{21}{75} = \frac{7}{25} \text{ radian}$$

### 15. Question

The radius of a circle is 30 cm. Find the length of an arc of this circle, if the length of the chord of the arc is 30 cm.

### Answer

Let AB be chord and O be the centre of the circle.

Here  $AO = BO = AB = 30$  cm

$\therefore \triangle AOB$  is an equilateral triangle.

Given radius = 30 cm

And  $\theta = 60^\circ$

We know that  $180^\circ = \pi \text{ rad} \Rightarrow 1^\circ = \pi/180 \text{ rad}$

$$\Rightarrow 60^\circ = \left(60 \times \frac{\pi}{180}\right) = \frac{\pi}{3} \text{ radian}$$

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \frac{\pi}{3} = \frac{\text{arc}}{30}$$

$$\therefore \text{arc} = \frac{30\pi}{3}$$

$$= 10\pi \text{ cm}$$

### 16. Question

A railway train is traveling on a circular curve of 1500 meters radius at the rate of 66 km/hr. Through what angle has it turned in 10 seconds?

### Answer

Given time is 10 seconds.

$$\text{And speed} = 66 \text{ km/h} = \frac{66 \times 1000}{3600} \text{ m/s}$$

We know that  $\text{speed} = \frac{\text{distance}}{\text{time}}$

$$\Rightarrow \frac{66 \times 1000}{3600} = \frac{\text{distance}}{\text{time}}$$

$$\Rightarrow \text{Distance} = \frac{66 \times 1000}{3600} \times 10 = \frac{1100}{6} \text{ m}$$

Now radius of curve = 1500 m

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \theta = \frac{\frac{1100}{6}}{1500}$$

$$= \frac{1100}{1500 \times 6} = \frac{11}{90} \text{ radian}$$

So, the train will turn  $11/90$  radian in 10 seconds.

### 17. Question

Find the distance from the eye at which a coin of 2 cm diameter should be held so as to conceal the full moon whose angular diameter is  $31'$ .

### Answer

Let PQ be the diameter of the coin and E be the eye of the observer.

Let the coin be kept at a distance  $r$  from the eye of the observer to hide the moon completely.

$$\Rightarrow \theta = 1' = \left(\frac{1}{60}\right)^\circ = \left(\frac{1}{60} \times \frac{\pi}{180}\right) \text{ rad}$$

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \frac{31}{60} \times \frac{\pi}{180} = \frac{2}{\text{radius}}$$

$$\therefore \text{radius} = \frac{180 \times 60 \times 2 \times 7}{31 \times 22}$$

$$= 221.7 \text{ cm}$$

### 18. Question

Find the diameter of the Sun in km supposing that it subtends an angle of  $32'$  at the eye of an observer. Given that the distance of the Sun is  $91 \times 10^6$  km.

### Answer

Let PQ be the diameter of the sun and E be the eye of the observer.

The distance between the Sun and Earth is large, so we will take PQ as arc PQ.

Given radius =  $91 \times 10^6$  km

$$\Rightarrow \theta = 32' = \left(\frac{32}{60}\right)^\circ = \left(\frac{32}{60} \times \frac{\pi}{180}\right) \text{ rad}$$

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \frac{32}{60} \times \frac{\pi}{180} = \frac{d}{91 \times 10^6}$$

$$\therefore d = \frac{32 \times 91 \times 10^6 \times 22}{60 \times 180 \times 7}$$

$$= 847407.4 \text{ km}$$

### 19. Question

If the arcs of the same length in two circles subtend angles  $65^\circ$  and  $110^\circ$  at the center, find the ratio of their radii.

### Answer

Let the angles subtended at the centers by the arcs and radii of first and second circles be  $\theta_1$  and  $r_1$  and  $\theta_2$  and  $r_2$ .

Then,

We know that  $180^\circ = \pi \text{ rad} \Rightarrow 1^\circ = \pi/180 \text{ rad}$

$$\Rightarrow \theta_1 = 65^\circ = \left(65 \times \frac{\pi}{180}\right) \text{ radian}$$

$$\Rightarrow \theta_2 = 110^\circ = \left(110 \times \frac{\pi}{180}\right) \text{ radian}$$

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \theta_1 = \frac{l}{r_1} \text{ and } \theta_2 = \frac{l}{r_2}$$

$$\Rightarrow r_1 = \frac{l}{\left(65 \times \frac{\pi}{180}\right)} \text{ and } r_2 = \frac{l}{\left(110 \times \frac{\pi}{180}\right)}$$

$$\therefore \frac{r_1}{r_2} = \frac{\frac{l}{\left(65 \times \frac{\pi}{180}\right)}}{\frac{l}{\left(110 \times \frac{\pi}{180}\right)}}$$

$$= \frac{110}{65}$$

$$= \frac{22}{13}$$

$$\therefore r_1:r_2 = 22:13$$

### 20. Question

Find the degree measure of the angle subtended at the centre of a circle of radius 100 cm by an arc of length 22 cm (Use  $\pi = \frac{22}{7}$ ).

### Answer

Given length of arc = 22 cm

And radius = 100 cm

We know that  $\theta = \frac{\text{arc}}{\text{radius}}$

$$\Rightarrow \theta = \frac{22}{100}$$

$$= \frac{11}{50} \text{radian}$$

$\therefore$  The angle subtended at the centre by the arc:

$$\Rightarrow \left(\frac{32}{60} \times \frac{\pi}{180}\right)^\circ = \left(\frac{11}{5} \times \frac{18}{220} \times 7\right)^\circ$$

$$= \left(\frac{63}{5}\right)^\circ$$

$$= 12^\circ 36'$$

## MCQ

### 1. Question

Mark the correct alternative in the following:

If D, G and R denote respectively the number of degrees, grades and radians in an angle, then

A.  $\frac{D}{100} = \frac{G}{90} = \frac{2R}{\pi}$

B.  $\frac{D}{90} = \frac{G}{100} = \frac{R}{\pi}$

C.  $\frac{D}{90} = \frac{G}{100} = \frac{2R}{\pi}$

D.  $\frac{D}{90} = \frac{G}{100} = \frac{R}{2\pi}$

### Answer

Let  $\theta$  be the angle which is measure in degree, radian and grade

We know that  $90^\circ = 1$  right angle

$$\Rightarrow 1^\circ = \frac{1}{90} \text{ right angle}$$

$$\Rightarrow D^\circ = \frac{D}{90} \text{ right angles}$$

$$\Rightarrow \theta = \frac{D}{90} \text{ right angle .....(1)}$$

Also we know that  $\pi$  radians = 2 right angles

$$\Rightarrow 1^c = \frac{2}{\pi} \text{ right angle}$$

$$\Rightarrow R = \frac{2}{\pi} \times R \text{ right angles}$$

$$\Rightarrow \theta = \frac{2}{\pi} \times R \text{ right angles .....(2)}$$

Also we know that, 100 grades = 1 right angle

$$\Rightarrow 1 \text{ grade} = \frac{1}{100} \text{ right angle}$$

$$\Rightarrow G \text{ grade} = \frac{G}{100} \text{ right angles}$$

$$\Rightarrow \theta = \frac{G}{100} \text{ right angles .....(3)}$$

From (1),(2) and (3)

$$\frac{D}{90} = \frac{2R}{\pi}$$

$$= \frac{G}{100}$$

## 2. Question

Mark the correct alternative in the following:

If the angles of a triangle are in A.P., then the measure of one of the angles in radians is

A.  $\frac{\pi}{6}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{2}$

D.  $\frac{2\pi}{3}$

## Answer

Here, angles of triangle are in A.P.

So, Let angles of triangle are  $a, a+d, a+2d$ .

We know that, sum of angles of triangle is  $\pi$ .

$$\therefore a+a+d+a+2d=\pi$$

$$\therefore 3a+3d=\pi$$

$$\therefore 3(a+d)=\pi$$

$$\therefore a+d = \frac{\pi}{3}$$

Also, by our assumption,  $a+d$  is one angle of triangle.

So, required measure of one of the angles is  $\frac{\pi}{3}$ .

## 3. Question

Mark the correct alternative in the following:

The angle between the minute and hour hands of a clock at 8:30 is

A.  $80^\circ$

B.  $75^\circ$

C.  $60^\circ$

D.  $105^\circ$

**Answer**

We know, in clock 1 rotation gives  $360^\circ$

i.e. 60 minutes =  $360^\circ$  and 12 hours =  $360^\circ$

So, 1 minute =  $6^\circ$  and 1 hour =  $30^\circ$

Now, For hour hand:

8 hours =  $8 \times 30^\circ = 240^\circ$  and for another 30 minute (which is half of hour) =  $30^\circ \div 2 = 15^\circ$

i.e. angle traced by hour hand is  $240^\circ + 15^\circ = 255^\circ$

Now, For minute hand:

30 minute =  $30 \times 6^\circ = 180^\circ$

i.e. angle traced by minute hand is  $180^\circ$ .

So, the angle between hour hand and minute hand =  $255^\circ - 180^\circ$

=  $75^\circ$

**4. Question**

Mark the correct alternative in the following:

At 3:40, the hour and minute hands of a clock are inclined at

A.  $\frac{2\pi^c}{3}$

B.  $\frac{7\pi^c}{12}$

C.  $\frac{13\pi^c}{18}$

D.  $\frac{3\pi^c}{4}$

**Answer**

We know, in clock 1 rotation gives  $360^\circ$

i.e. 60 minutes =  $360^\circ$  and 12 hours =  $360^\circ$

So, 1 minute =  $6^\circ$  and 1 hour =  $30^\circ$

Now, For hour hand:

3 hours =  $3 \times 30^\circ = 90^\circ$  and for another 40 minute =  $(30^\circ \div 60) \times 40 = 20^\circ$

i.e. angle traced by hour hand is  $90^\circ + 20^\circ = 110^\circ$

Now, for minute hand:

40 minute =  $40 \times 6^\circ = 240^\circ$

i.e. angle traced by minute hand is  $240^\circ$ .

So, the angle between hour hand and minute hand =  $240^\circ - 110^\circ$

=  $130^\circ$

$$= 130^\circ \times \frac{\pi^c}{180}$$

$$= \frac{13\pi^c}{18}$$

### 5. Question

Mark the correct alternative in the following:

If the arcs of the same length in two circles subtend angles  $65^\circ$  and  $110^\circ$  at the centre, then the ratio of the radii of the circles is

- A. 22 : 13
- B. 11 : 13
- C. 22 : 15
- D. 21 : 13

### Answer

Let radius of two circles be the  $r_1$  and  $r_2$

Let  $\theta_1$  and  $\theta_2$  be the subtend angles of arcs of two circles

i.e.  $\theta_1=65^\circ$  and  $\theta_2=110^\circ$

We know that arc length,

$$l = r \times \theta$$

Here, arc lengths of two circles are same.

$$\therefore r_1 \times \theta_1 = r_2 \times \theta_2$$

$$\therefore \frac{r_1}{r_2} = \frac{\theta_2}{\theta_1} = \frac{110}{65}$$

$$\therefore \frac{r_1}{r_2} = \frac{11 \times 2}{13}$$

$$\therefore r_1 : r_2 = 22 : 13$$

### 6. Question

Mark the correct alternative in the following:

If OP makes 4 revolutions in one second, the angular velocity in radians per second is

- A.  $\pi$
- B.  $2\pi$
- C.  $4\pi$
- D.  $8\pi$

### Answer

We know that, 1 revolution =  $2 \times \pi$  radians

Now, Angular velocity =  $\frac{\text{Distance}}{\text{Time}}$

$$= \frac{4 \text{ revolutions}}{1 \text{ second}}$$

$$= \frac{4 \times 2 \times \pi}{1}$$

$$= 8 \times \pi$$

### 7. Question



Mark the correct alternative in the following:

A circular wire of radius 7 cm is cut and bent again into an arc of a circle of radius 12 cm. The angle subtended by the arc at the centre is

- A.  $50^\circ$
- B.  $210^\circ$
- C.  $100^\circ$
- D.  $60^\circ$
- E.  $195^\circ$

**Answer**

Here, radius of circular wire is  $r=7$  cm

So, length of wire  $=2 \times \pi \times r$

$$=2 \times \pi \times 7$$

$$=14 \times \pi$$

Wire is cut and bent again into an arc of a circle of radius 12 cm.

So, length of arc=length of wire  $=14 \times \pi$

We know, angle subtended by the arc is given by,

$$\theta = \frac{\text{length of arc}}{\text{radius}}$$

$$= \frac{14\pi}{12}$$

$$= \frac{14\pi}{12} \times \frac{180^\circ}{\pi}$$

$$=210^\circ$$

**8. Question**

Mark the correct alternative in the following:

The radius of the circle whose arc of length  $15\pi$  cm makes an angle of  $3\pi/4$  radian at the centre is

- A. 10 cm
- B. 20 cm
- C.  $11\frac{1}{4}$  cm
- D.  $22\frac{1}{2}$  cm

**Answer**

Here, arc length  $l=15\pi$  cm

$$\text{Angle } \theta = \frac{3\pi}{4}$$

We know, angle subtended by the arc is given by,

$$\theta = \frac{\text{length of arc}}{\text{radius}}$$

$$\therefore \text{radius} = \frac{1}{\theta}$$

$$= \frac{15\pi}{3\pi} \times 4$$

$$= 20 \text{ cm}$$

[Bodhiyla.com](http://Bodhiyla.com)